

A-Potene™

DESCRIPTION

A-Potene™, provided by Douglas Laboratories, contains 25,000 IU of beta-carotene in each softgel.

FUNCTIONS

Beta-carotene is a safe precursor of vitamin A. Human intestinal enzymes convert dietary beta-carotene into active vitamin A. The extent of this conversion appears to be based on the need for vitamin A since even very large amounts of beta-carotene do not increase plasma or liver vitamin A levels above normal. One milligram of beta-carotene provides 1,667 I.U. of vitamin A activity. Beta-carotene not converted to vitamin A is absorbed intact and has other functions in the body. Beta-carotene is an important antioxidant nutrient that protects healthy cells from oxidative and free radical damage. Free radicals are unstable chemicals formed in the body during metabolism and from exposure to environmental sources such as pollution and cigarette smoke. Free radicals are necessary for energy metabolism and immune function, but when an excessive number of free radicals are formed, they can attack healthy cells, especially their membrane lipids and proteins. This, in turn, is thought to contribute to a number of degenerative diseases. Beta-carotene efficiently quenches a chemically reactive species of oxygen called singlet oxygen. Singlet oxygen participates in oxidative reactions which can impair or destroy important cellular components, such as membranes, nucleic acids (DNA), and enzymes. In addition, singlet oxygen reactions can generate free radicals which lead to further damage. Beta-carotene's long, conjugated molecule can absorb the energy of singlet oxygen and release it safely as heat. In the process, singlet oxygen is converted to less dangerous, regular oxygen. One molecule of beta-carotene can quench about a thousand molecules of singlet oxygen. Numerous studies have demonstrated the importance of beta-carotene for normal immune function. The immune benefits of beta-carotene are due to its vitamin A precursor role and its antioxidant functions.

Beta-carotene is present in selected fruits and vegetables such as carrots, sweet potatoes, and green leafy vegetables. The U.S. Department of Agriculture and the National Cancer Institute recommend an intake of 5 mg (8,335 I.U.) or more of beta-carotene. Unfortunately, most diets provide only 1-2 mg (1,667 - 3,334 I.U.) per day.

INDICATIONS

A-Potene™ may be a useful dietary supplement for those who wish to increase their daily beta-carotene intake.

FORMULA (#7993)

Each Softgel Contains:

Vitamin A (as 100% Beta-Carotene) 25,000 IU

SUGGESTED USE

Adults take 1 softgel daily or as directed by healthcare professional.

SIDE EFFECTS

No adverse side effects have been reported.

STORAGE

Store in a cool, dry place, away from direct light. Keep out of reach of children.

A-Potene™

REFERENCES

- Agarwal S, Rao AV. Carotenoids and chronic diseases. *Drug Metabol Drug Interact* 2000;17(1-4):189-210
2000. PMID:15130.
- Allard JP, Royall D, Kurian R, Muggli R, Jeejeebhoy KN. Effects of beta-carotene supplementation on lipid peroxidation in humans. *Am J Clin Nutr* 1994;59:884-90.
- Baron JA, Cole BF, Mott L, et al. Neoplastic and antineoplastic effects of beta-carotene on colorectal adenoma recurrence: results of a randomized trial. *J Natl Cancer Inst* 2003;95(10):717-722.
- Bendich A. Beta-carotene and the immune response. *Proc Nutr Soc* 1991;50:263-74.
- Bertram JS. Carotenoids and gene regulation. *Nutr Rev* 1999;57:182-91.
- Buring JE, Hennekens CH. beta-carotene and cancer chemoprevention. *J Cell Biochem Suppl* 1995;22:226-30.
- Christen WG, Manson JE, Glynn RJ, et al. Beta carotene supplementation and age-related maculopathy in a randomized trial of US physicians. *Arch Ophthalmol* 2007 Mar;125(3):333-9.
- Comstock GW, Burke AE, Hoffman SC, Helzlsouer KJ, Bendich A, Masi AT, Norkus EP, Malamet RL, Gershwin ME. Serum concentrations of alpha tocopherol, beta carotene, and retinol preceding the diagnosis of rheumatoid arthritis and systemic lupus erythematosus. *Ann Rheum Dis* 1997;56:323-5.
- Cooper DA, Eldridge AL, Peters JC. Dietary carotenoids and certain cancers, heart disease, and age-related macular degeneration: a review of recent research. *Nutr Rev* 1999;57:201-14.
- Daudu PA, Kelley DS, Taylor PC, Burri BJ, Wu MM. Effect of a low beta-carotene diet on the immune functions of adult women. *Am J Clin Nutr* 1994;60:969-72.
- Gey KF, Moser UK, Jordan P, Stahelin HB, Eichholzer M, Ludin E. Increased risk of cardiovascular disease at suboptimal plasma concentrations of essential antioxidants: an epidemiological update with special attention to carotene and vitamin C. *Am J Clin Nutr* 1993;57:787S-797S.
- Grodstein F, Kang JH, Glynn RJ, et al. A randomized trial of beta carotene supplementation and cognitive function in men: the Physicians' Health Study II. *Arch Intern Med* 2007 Nov 12;167(20):2184-90.
- Hennekens CH, Buring JE, Manson JE, Stampfer M, Rosner B, Cook NR, Belanger C, LaMotte F, Gaziano JM, Ridker PM, Willett W, Peto R. Lack of effect of long-term supplementation with beta carotene on the incidence of malignant neoplasms and cardiovascular disease [see comments]. *N Engl J Med* 1996;334:1145-9.
- Herraiz LA, Hsieh WC, Parker RS, Swanson JE, Bendich A, Roe DA. Effect of UV exposure and beta-carotene supplementation on delayed type hypersensitivity response in healthy older men. *J Am Coll Nutr* 1998;17:617-24.
- Kornhauser A, Wamer WG, Lambert LA, Wei RR. beta-Carotene inhibition of chemically induced toxicity in vivo and in vitro. *Food Chem Toxicol* 1994;32:149-54.
- Krinsky NI. Carotenoids as antioxidants. *Nutrition* 2001 Oct;17(10):815-7 2001. PMID:15110.
- Lambert LA, Wamer WG, Wei RR, Lavu S, Chirtel SJ, Kornhauser A. The protective but nonsynergistic effect of dietary beta-carotene and vitamin E on skin tumorigenesis in Skh mice. *Nutr Cancer* 1994;21:1-12.
- Levy Y, Zaltsberg H, Ben-Amotz A, Kanter Y, Aviram M. Dietary supplementation of a natural isomer mixture of beta-carotene inhibits oxidation of LDL derived from patients with diabetes mellitus. *Ann Nutr Metab* 2000;44(2):54-60.
- Murata T, Tamai H, Morinobu T, Manago M, Takenaka H, Hayashi K, Mino M. Effect of long-term administration of beta-carotene on lymphocyte subsets in humans. *Am J Clin Nutr* 1994;60:597-602.
- Nishino H. Cancer chemoprevention by natural carotenoids and their related compounds. *J Cell Biochem Suppl* 1995;22:231-5.

A-Potene™

Noonan FP, Webber LJ, De Fabo EC, Hoffman HA, Bendich A, Mathews-Roth M. Dietary beta-carotene and ultraviolet-induced immunosuppression. *Clin Exp Immunol* 1996;103:54-60.

Ozhogina OA, Kasaikina OT. Beta-carotene as an interceptor of free radicals. *Free Radic Biol Med* 1995;19:575-81.

Santos MS, Leka LS, Ribaya-Mercado JD, Russell RM, Meydani M, Hennekens CH, Gaziano JM, Meydani SN. Short- and long-term betacarotene supplementation do not influence T cell-mediated immunity in healthy elderly persons. *Am J Clin Nutr* 1997;66:917-24.

Santos MS, Meydani SN, Leka L, Wu D, Fotouhi N, Meydani M, Hennekens CH, Gaziano JM. Natural killer cell activity in elderly men

is enhanced by beta- carotene supplementation. *Am J Clin Nutr* 1996;64:772-7.

Teikari JM, Virtamo J, Rautalahti M, Palmgren J, Liesto K, Heinonen

OP. Long-term supplementation with alpha-tocopherol and betacarotene and age-related cataract. *Acta Ophthalmol Scand*

1997;75:634-40.

Wamer WG, Wei RR, Matusik JE, Kornhauser A, Dunkel VC. beta-Carotene uptake, metabolism, and distribution in vitro. *Ann N Y Acad Sci* 1993;691:284-6.

Wei RR, Wamer WG, Lambert LA, Kornhauser A. beta-Carotene uptake and effects on intracellular levels of retinol in vitro. *Nutr Cancer* 1998;30:53-8.

Young AJ, Lowe GM. Antioxidant and prooxidant properties of carotenoids. *Arch Biochem Biophys* 2001 Jan 1;385(1):20-7 2001. PMID:15120

For more information on A-Potene™ visit douglaslabs.com

† These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

Manufactured by
Douglas Laboratories
600 Boyce Road
Pittsburgh, PA 15205
800-245-4440
douglaslabs.com



**You trust Douglas Laboratories.
Your patients trust you.**