

201910-120

Dietary Supplement

**Wobenzym® plus**

**Cautions and Warnings:** If you have gastrointestinal lesions /ulcers, bleeding disorders, or are taking anticoagulant / blood thinners, anti-inflammatory or antibiotic medications, or are about to have surgery, consult a healthcare practitioner prior to use. Do not use if you are pregnant or breast-feeding. If you experience nausea, vomiting or hypersensitivity / allergic reaction while taking Wobenzym, discontinue use and consult your healthcare practitioner.

**Recommended dose:**  
Adults: Take 2 tablets twice daily at least 45 minutes before meals with water.

**Store in a cool dry place.**  
**Do not use if safety seal is broken or missing.** Keep out of reach of children.

**Distributed by:**  
Mucos USA  
112 Technology Drive  
Pittsburgh, PA 15108  
Wobenzym.com

Helps to maintain healthy joints, mobility and flexibility<sup>1</sup>

Temporary relief from everyday aches, pains and muscle soreness<sup>1</sup>

Speedy recovery following exertion<sup>1</sup>



120 enteric-coated tablets

1 These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

	Amount Per Serving	% Daily Value
Bromelain** (from pineapple ( <i>Ananas comosus</i> var <i>comosus</i> ) stem)	270 mg (1350 FIP*** -units)	*
Trypsin (from porcine ( <i>Sus scrofa</i> ) or bovine ( <i>Bos taurus</i> ) pancreas)	144 mg (4320 FIP*** -units)	*
Rutoside trihydrate (Rutin) (from <i>Staphylocarpus japonicum</i> flower bud)	300 mg	*

**Other ingredients:** Microcrystalline cellulose, calcium phosphate, hydroxypropyl cellulose, vegetable stearate, enteric coating (sodium alginate, medium chain triglycerides, ethylcellulose, stearic acid, oleic acid, purified water) purified water.

201910-240

Dietary Supplement

**Wobenzym® plus**

**Cautions and Warnings:** If you have gastrointestinal lesions /ulcers, bleeding disorders, or are taking anticoagulant / blood thinners, anti-inflammatory or antibiotic medications, or are about to have surgery, consult a healthcare practitioner prior to use. Do not use if you are pregnant or breast-feeding. If you experience nausea, vomiting or hypersensitivity / allergic reaction while taking Wobenzym, discontinue use and consult your healthcare practitioner.

**Recommended dose:**  
Adults: Take 2 tablets twice daily at least 45 minutes before meals with water.

**Store in a cool dry place.**  
**Do not use if safety seal is broken or missing.** Keep out of reach of children.

**Distributed by:**  
Mucos USA  
112 Technology Drive  
Pittsburgh, PA 15108  
Wobenzym.com

Helps to maintain healthy joints, mobility and flexibility<sup>1</sup>

Temporary relief from everyday aches, pains and muscle soreness<sup>1</sup>

Speedy recovery following exertion<sup>1</sup>



240 enteric-coated tablets

1 These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

	Amount Per Serving	% Daily Value
Bromelain** (from pineapple ( <i>Ananas comosus</i> var <i>comosus</i> ) stem)	270 mg (1350 FIP*** -units)	*
Trypsin (from porcine ( <i>Sus scrofa</i> ) or bovine ( <i>Bos taurus</i> ) pancreas)	144 mg (4320 FIP*** -units)	*
Rutoside trihydrate (Rutin) (from <i>Staphylocarpus japonicum</i> flower bud)	300 mg	*

**Other ingredients:** Microcrystalline cellulose, calcium phosphate, hydroxypropyl cellulose, vegetable stearate, enteric coating (sodium alginate, medium chain triglycerides, ethylcellulose, stearic acid, oleic acid, purified water) purified water.

201910-480

Dietary Supplement

**Wobenzym® plus**

**Cautions and Warnings:** If you have gastrointestinal lesions /ulcers, bleeding disorders, or are taking anticoagulant / blood thinners, anti-inflammatory or antibiotic medications, or are about to have surgery, consult a healthcare practitioner prior to use. Do not use if you are pregnant or breast-feeding. If you experience nausea, vomiting or hypersensitivity / allergic reaction while taking Wobenzym, discontinue use and consult your healthcare practitioner.

**Recommended dose:**  
Adults: Take 2 tablets twice daily at least 45 minutes before meals with water.

**Store in a cool dry place.**  
**Do not use if safety seal is broken or missing.** Keep out of reach of children.

**Distributed by:**  
Mucos USA  
112 Technology Drive  
Pittsburgh, PA 15108  
Wobenzym.com

Helps to maintain healthy joints, mobility and flexibility<sup>1</sup>

Temporary relief from everyday aches, pains and muscle soreness<sup>1</sup>

Speedy recovery following exertion<sup>1</sup>



480 enteric-coated tablets

1 These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

	Amount Per Serving	% Daily Value
Bromelain** (from pineapple ( <i>Ananas comosus</i> var <i>comosus</i> ) stem)	270 mg (1350 FIP*** -units)	*
Trypsin (from porcine ( <i>Sus scrofa</i> ) or bovine ( <i>Bos taurus</i> ) pancreas)	144 mg (4320 FIP*** -units)	*
Rutoside trihydrate (Rutin) (from <i>Staphylocarpus japonicum</i> flower bud)	300 mg	*

**Other ingredients:** Microcrystalline cellulose, calcium phosphate, hydroxypropyl cellulose, vegetable stearate, enteric coating (sodium alginate, medium chain triglycerides, ethylcellulose, stearic acid, oleic acid, purified water) purified water.