

EPA/GLA Forte
Cardiovascular Health

DESCRIPTION

EPA/GLA Forte softgels, provided by Douglas Laboratories®, contain a marine lipid concentrate processed by molecular distillation and borage seed oil. EPA/GLA Forte is an excellent source of both omega-3 fatty acids, providing eicosapentaenoic acid (EPA, 18%) and docosahexaenoic acid (DHA, 12%) in their natural triglyceride form, as well as the omega-6 fatty acid gamma linolenic acid (GLA).

FUNCTIONS

Typical diets in developed countries deliver large amounts of saturated fatty acids and the polyunsaturated omega-6 linoleic and arachidonic acids and low levels of omega-3 fatty acids. Throughout evolution, humans were accustomed to diets providing roughly equal amounts of omega-6 and omega-3 fatty acids. However, during the last 200 years, the ratio of dietary omega-6:omega-3 fatty acids increased from about 1:1 to 20-25:1. Leading health professionals now recommend ratios between 4:1 and 10:1. A healthy balance of dietary omega-6 and omega-3 fatty acids appears to be a prerequisite for normal immune function. Dietary linoleic acid (18:2 omega-6) is a precursor to arachidonic acid (20:4) which in turn is a precursor for pro-inflammatory prostaglandin E2 and leukotriene B4, and platelet aggregating thromboxane A2. Although GLA is a precursor of arachidonic acid, it also competes with arachidonic acid effectively, and may help downregulate the formation of excessive levels of pro-inflammatory 2-series prostaglandins and other immune mediators. The omega-3 fatty acids EPA and DHA provide a natural counterbalance to the effects of excess omega-6 fatty acids, because they serve as precursors for the anti-inflammatory prostaglandins E1 and E3, and decrease the formation of prostaglandin E2 and thromboxane A2. Thus, a balanced dietary intake of all of these fatty acids appears to be necessary to maintain a healthy immune response during normal inflammatory processes. The omega-3 fatty acids also have important functions for visual acuity and possibly in maintaining normal blood lipoprotein levels and the healthy metabolism of cholesterol.

INDICATIONS

EPA/GLA Forte softgels may be a useful dietary supplement for those who do not consume enough oily fish and other sources of omega-3 fatty acids to receive desirable amounts of healthy long chain omega-3 fatty acids. EPA/GLA Forte may also be a useful dietary adjunct for those who wish to supplement their diets with gamma-linolenic acid.

FORMULA (#7040)

Each Softgel Capsule Contains:

Natural Triglyceride Marine Lipid Concentrate	667 mg
Borage Oil	62.5 mg
Vitamin E (d-alpha Tocopherol with Mixed Tocopherols)	10 I.U.
Providing:	
Pure Eicosapentaenoic Acid (EPA)	120 mg
Gamma Linolenic Acid (GLA)	15 mg

SUGGESTED USE

Adults take 1 to 4 softgels 1 to 3 times daily or as directed by physician.

SIDE EFFECTS

No adverse side effects have been reported.

EPA/GLA Forte
Cardiovascular Health
STORAGE

Store in a cool, dry place, away from direct light. Keep out of reach of children.

REFERENCES

Albert CM, Hennekens CH, O'Donnell CJ, et al. Fish consumption and risk of sudden cardiac death [see comments]. *Jama* 1998;279:23-8.

Cerbone AM, Cirillo F, Coppola A, et al. Persistent impairment of platelet aggregation following cessation of a short-course dietary supplementation of moderate amounts of N-3 fatty acid ethyl esters. *Thromb Haemost* 1999;82:128-33.

Christensen JH, Gustenhoff P, Korup E, et al. [n-3 polyunsaturated fatty acids, heart rate variability and ventricular arrhythmias in post-AMI-patients. A clinical controlled trial]. *Ugeskr Laeger* 1997;159:5525-9.

Christensen JH, Korup E, Aaroe J, et al. Fish consumption, n-3 fatty acids in cell membranes, and heart rate variability in survivors of myocardial infarction with left ventricular dysfunction. *Am J Cardiol* 1997;79:1670-3.

de Deckere EA, Korver O, Verschuren PM, et al. Health aspects of fish and n-3 polyunsaturated fatty acids from plant and marine origin. *Eur J Clin Nutr* 1998;52:749-53.

Fan YY, Chapkin RS. Importance of dietary gamma-linolenic acid in human health and nutrition. *J Nutr* 1998;128:1411-4.

Horrocks LA, Yeo YK. Health benefits of docosahexaenoic acid (dha). *Pharmacol Res* 1999;40:211-25.

Leng GC, Lee AJ, Fowkes FG, et al. Randomized controlled trial of gamma-linolenic acid and eicosapentaenoic acid in peripheral arterial disease. *Clin Nutr* 1998;17:265-71.

Simopoulos AP. Essential fatty acids in health and chronic disease. *Am J Clin Nutr* 1999;70:560S-569S.

Takahata K, Monobe K, Tada M, et al. The benefits and risks of n-3 polyunsaturated fatty acids. *Biosci Biotechnol Biochem* 1998;62:2079-85.

Uauy R, Mena P, Valenzuela A. Essential fatty acids as determinants of lipid requirements in infants, children and adults. *Eur J Clin Nutr* 1999;53 Suppl 1:S66-77.

For more information on EPA/GLA visit douglaslabs.com

† These statements have not been evaluated by the Food and Drug Administration.
This product is not intended to diagnose, treat, cure, or prevent any disease.

Manufactured by
Douglas Laboratories
600 Boyce Road
Pittsburgh, PA 15205
800-245-4440
douglaslabs.com



**You trust Douglas Laboratories.
Your patients trust you.**